

# L-Carnitine Tartrate SAP

Science-based fatty acid metabolism and antioxidant support

L-Carnitine plays an important role in fatty acid metabolism, and has important antioxidant and anti-inflammatory properties.<sup>[1]</sup> L-Carnitine is a nonessential amino acid synthesized primarily in the liver and kidneys from the amino acids lysine and methionine. L-Carnitine is stored in skeletal muscles, brain, heart, and sperm, and therefore plays a functional role in each of these tissues. Deficiency in L-carnitine may be associated with certain medications, angina, and vegan or vegetarian diets. Symptoms of deficiency may include fatigue, muscle weakness, and a decreased tolerance to metabolic stress.<sup>[1]</sup>

## ACTIVE INGREDIENTS

Each capsule of L-Carnitine Tartrate SAP contains:

L-Carnitine ..... 500 mg  
From 750 mg of L-carnitine tartrate.

**This product is non-GMO.**

**Contains no:** Gluten, soy, wheat, corn, eggs, dairy, yeast, citrus, preservatives, artificial flavor or color, starch, or sugar.

**L-Carnitine Tartrate SAP** contains 90 capsules

## DIRECTIONS FOR USE

**Adults: Start with 1 capsule twice daily with food** and gradually increase to 2 capsules twice daily with food or as directed by your healthcare practitioner. Take 2–4 hours prior to exercise. Consult a healthcare practitioner for use beyond 6 months.

## INDICATIONS

**L-Carnitine Tartrate SAP** may:

- Be used to help prevent stable angina.
- Be used to help prevent pain associated with peripheral vascular disease (PVD).
- Increase sperm count and mobility.
- Help prevent cancer cachexia.
- Improve recovery time for athletes.

## SAFETY AND INTERACTIONS

- L-Carnitine is considered to be a safe supplement with rare side effects that may include nausea or gastric upset. L-Carnitine does interact with some medications, so if you are taking any of the medications below, please speak with your healthcare practitioner before taking L-carnitine:
  - Isotretinoin (Accutane) can cause side effects similar to those seen with carnitine deficiency, including muscle pain and weakness, high cholesterol, and liver concerns. Taking L-carnitine with this medication may improve these side effects.
  - Valproic acid (Depakote) – Taking L-carnitine concurrently with this medication may prevent any deficiency and reduce the side effects of valproic acid.
  - Doxorubicin – Concurrent use of L-carnitine may help protect cardiac cells against the toxicity that can result from doxorubicin.
  - Thyroid hormone – L-Carnitine may reduce the amount of thyroid hormone that is able to get into cells.

## PURITY, CLEANLINESS, AND STABILITY

All ingredients listed for all **L-Carnitine Tartrate SAP** lot numbers have been tested by a third-party laboratory for identity, potency, and purity.

Scientific Advisory Panel (SAP):  
adding nutraceutical research  
to achieve optimum health



351, Rue Joseph-Carrier, Vaudreuil-Dorion, Quebec, J7V 5V5  
T 1 866 510 3123 • F 1 866 510 3130 • nfh.ca

## L-CARNITINE BIOCHEMISTRY

Carnitine is a cofactor that is required to metabolize free long-chain fatty acids into acylcarnitines, which are then transported into mitochondria for  $\beta$ -oxidation.<sup>[2]</sup> This form of energy production is the primary fuel source for both heart and skeletal muscle. Carnitine can be synthesized via methylated L-lysine with S-adenosyl methionine (SAM).<sup>[2]</sup> Cofactors important for this process include magnesium, ascorbic acid, iron, pyridoxal-5'-phosphate, niacin, and methionine, as well as the cofactors necessary for the methylation pathway.<sup>[2]</sup> If patients are deficient in any of the cofactors, this may result in a relative deficiency of carnitine, which may have an impact on proper functioning of skeletal muscle, heart, or sperm function.

## ANGINA

L-Carnitine may help improve duration of exercise and recovery time in patients with stable angina.<sup>[2]</sup> A six-month trial using 2 g/d L-carnitine in patients with exercise-induced stable angina demonstrated a reduction in the number of premature ventricular contractions at rest, improved exercise tolerance, increased maximal systolic arterial blood pressure, and reduced ST-segment depression during maximal effort.<sup>[2]</sup> There was also a reduction in the number of cardioactive pharmaceuticals needed in patients supplementing L-carnitine.<sup>[2]</sup>

In a separate randomized, placebo-controlled, crossover study investigating stable effort-induced angina, patients were given L-carnitine or placebo.<sup>[2]</sup> Twenty-two percent of the patients in the L-carnitine group reported being free of angina, compared to nine percent in the placebo group.<sup>[2]</sup> Results indicated a reduction in ECG indices of ischemia and improved exercise tolerance with L-carnitine supplementation.<sup>[2]</sup>

## INTERMITTENT CLAUDICATION

A review article exploring the use of supplementing L-carnitine in patients with intermittent claudication examined 17 articles that met their inclusion criteria.<sup>[3]</sup> Researchers found that of the 5 randomized control trials, 4 demonstrated significant improvements in walking performance following dosing of between 300 mg/d and 600 mg/d of oral L-carnitine or propionyl-L-carnitine (PLC).<sup>[3]</sup> Average improvements compared to placebo showed an improvement in pain-free walking distance as well as for maximal walking distance by 23–132 m and 104 m, respectively, following carnitine intervention.<sup>[3]</sup>

## MALE FERTILITY

L-Carnitine plays an essential role in maintaining male fertility. In a study examining sperm motility, count,

and morphology, researchers found that infertile subjects had a significantly lower amount of seminal free L-carnitine compared to fertile controls, and that subjects in the azoospermic group had the lowest levels of L-carnitine.<sup>[4]</sup> In another study, researchers found a significant positive correlation between seminal plasma total carnitine concentration and total sperm count.<sup>[5]</sup> Researchers concluded that seminal carnitine levels may be a useful test when evaluating male infertility.<sup>[5]</sup>

## CANCER CACHEXIA

In patients with cancer, cachexia is a major concern and a likely determinate of longevity. In a review study, researchers presented evidence regarding the use of carnitine supplementation for treating cachexia in patients with cancer. Data showed that cancer patients had low serum carnitine levels, and that L-carnitine supplementation resulted in improvement in fatigue as well as quality of life.<sup>[1]</sup>

## ATHLETES

L-Carnitine can enhance vascular endothelial function, which may lead to improving blood flow to muscle tissue, resulting in a decrease in hypoxic stress.<sup>[6]</sup> In a direct assessment of muscle-tissue damage using an MRI, researchers found that L-carnitine supplementation reduced muscle damage related to hypoxic stress.<sup>[6]</sup>

Other aspects of L-carnitine with regards to athletic performance have had mixed results. There are some studies that have shown a positive impact on  $VO_{2\max}$ ; however, others have not supported those findings.<sup>[2]</sup> There are also some studies that demonstrate using L-carnitine during exercise may enhance fat metabolism, while others fail to observe a benefit in body-fat percentage reduction.<sup>[2]</sup> It appears likely that if an athlete is deficient in L-carnitine, they may benefit from supplementation; however, athletes with sufficient L-carnitine supplementation may have limited effect on  $VO_{2\max}$  or fat metabolism.

## REFERENCES

1. Silvério, R., et al. "L-Carnitine and cancer cachexia: Clinical and experimental aspects." *Journal of Cachexia, Sarcopenia and Muscle* Vol. 2, No. 1 (2011): 37–44.
2. Kelly, G. "L-Carnitine: Therapeutic applications of a conditionally essential amino acid." *Alternative Medicine Review* Vol. 3, No. 5 (1998): 345–360.
3. Delaney, C.L., et al. "A systematic review to evaluate the effectiveness of carnitine supplementation in improving walking performance among individuals with intermittent claudication." *Atherosclerosis* Vol. 229, No. 1 (2013): 1–9.
4. Ahmed, S.D., et al. "Role of L-carnitine in male infertility." *Journal of the Pakistan Medical Association* Vol. 61, No. 8 (2011): 732–736.
5. Gürbüz, B., et al. "Relationship between semen quality and seminal plasma total carnitine in infertile men." *Journal of Obstetrics and Gynaecology* Vol. 23, No. 6 (2003): 653–656.
6. Huang, A. and K. Owen. "Role of supplementary L-carnitine in exercise and exercise recovery." *Medicine and Sport Science* Vol. 59 (2012): 135–142.